EIC Forward GEM Tracking using PHGenFit and PHG4Hit

Jin Huang(BNL),

Haiwang Yu (NMSU)

Jul 06, 2016

Outline

Motivation: Produce fsPHENIX tracking performance with Geant simulation + Kalman Filter.

- More realistic than Sagitta calculation.
- Serves as prototype for forward sPHENIX tracking software

Procedure:

- Simulate single muons using PHG4SimpleEventGenerator. $\Longrightarrow$ PHG4Hit
- Smear PHG4Hit from muon track according to given detector resolution.
- Use that smeared PHG4Hit and GenFit Kalman Filter do the fitting.

Contents:

- Method Verification: test with 3-layer vacuum detector.
- Test using EIC concept FGEM, arXiv:1402.1209
- Companion with previous studies.

Method Verification:
Using 3-layer vertical plane tracking

3-layer vertical plane testing setup

First, we tested the code using 3-layer vertical planes:

- The world and all detector material is G4_Galactic.
- 3-layer vertical planes with black holes outside.
- δr = 1cm
- r$\delta\phi$ = 100 μm



Vacuum, 3-layer, η = 3.0, r$\delta\phi$ = 100 μm, δr = 1cm

PHGenFit results are very similar with the Geant4 Sagitta calculation:

- Blue circle: $\sigma_{Sagitta}$/Sagitta vs. p from Geant4 simulation
- Black dot: σ_p/p vs. p from PHGenFit Kalman fitting.



EIC Concept FGEM tracker: Momentum resolution σ_p/p

5(4)-station Forward GEM Detector

- Detector setup using G4_FGEM_fsPHENIX.C in sPHENIX master branch.
- z = 17 cm (1.01 < η < 2.7), 62 cm (2.15 < η <4.0), 120, 160, 275 (1.45 < η < 4) cm
- δr = 1 cm, r$\delta\phi$ = 100 μm (η = 1.5~2.5) 50 μm (η = 2.5~4)
- sPHENIX field, no passive piston.
- **Tracking used vertex** from smeared truth vertex (0,0,0) and 50 μm resolution.





how to calculate σ_p/p

- 2D histo: $(p_{Reco} - p_{True})/p_{True}$ vs p_{True}, (right plot)
- For each slice of p_{True}, fit with Gaussian, extract mean as offset (Grey dots), sigma as momentum resolution (σ_p: Black dots), left plot.
- We also calculated $<\sigma_{Sagitta}/Sagitta>$ as a reference (Blue circles): Sagitta is calculated using vertex, station at 120cm and station at 275 cm.





Result: σ_p/p for different pseudo-rapidity



Details for each curve are in the backups.
PDF format: https://www.phenix.bnl.gov/WWW/p/draft/yuhw/fsPHENIX/FGEM_PHGenFit.pdf

Compare with previous studies

Compare with previous studies, Jin's MatLab calculation

Jin's calculation based on vertex + optimum Sagitta plane + 300cm last station.



GenFit fitting for η = 3.0, corresponding to magenta curve in left plot.

- The linear term, p1, from the σ_S/S **is consistent with left plot,** both are ~0.25%.
- The p1 term from full GenFit Kalman is better than σ_S/S . That could be caused by that we have more stations in full Kalman.



Compare with BeAST:

- Resolution is about x2 of the BeAST, despite BeAST use x2 stronger magnetic field + silicon tracker, as we used a much longer tracking arm.
- Our higher momentum resolution could be improved by also switching GEM to high precision silicon tracker.





sPHENIX FGEM



BeAST

Compare with previous studies, arXiv:1402.1209

- Linear term p1 is better than previous Sagitta tracking study by **using a vertex in the fitting**.
- Constant term p0, (Multiple Scattering Effect) is constant with previous Sagitta tracking study.





MS: < 1% (low η) to 3% (η = 3)

arXiv:1402.1209

Summary

- Tracking prototype made based on PHGenFit and PHG4Hit.
  - https://github.com/HaiwangYu/FGEM_PHGenFit_G4Hit_Tracking

- Produced performance plots for the EIC concept forward tracking detector.
  - Results consistent with previous studies.

Backups:

η = 1.5, r$\delta\phi$ = 100 μm





η = 2.0, r$\delta\phi$ = 100 μm





η = 2.5, r$\delta\phi$ = 100 μm





η = 3.0, r$\delta\phi$ = 50 μm





η = 3.5, r$\delta\phi$ = 50 μm